



## University of Groningen

### Education, diet, and incident cardiovascular disease

Jacobs, David R.; Kromhout, Daan

*Published in:*  
The Lancet Global Health

*DOI:*  
[10.1016/S2214-109X\(19\)30187-1](https://doi.org/10.1016/S2214-109X(19)30187-1)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2019

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Jacobs, D. R., & Kromhout, D. (2019). Education, diet, and incident cardiovascular disease: ecological interactions and conclusions. *The Lancet Global Health*, 7(6), E684-E685. [https://doi.org/10.1016/S2214-109X\(19\)30187-1](https://doi.org/10.1016/S2214-109X(19)30187-1)

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Education, diet, and incident cardiovascular disease: ecological interactions and conclusions



The Prospective Urban Rural Epidemiologic (PURE) study<sup>1</sup> is an impressive epidemiological undertaking, coordinating clinical examinations and multi-year follow-up across more than 18 countries with more than 180 000 participants. This study is a successor of the Seven Countries study,<sup>2,3</sup> which was led by Ancel Keys and was done across seven countries with more than 12 000 participants. Keys made the fundamental observation that coronary heart disease and, correspondingly, serum total cholesterol differ dramatically across cultures, and he strongly asserted that diet is important in between-country differences.<sup>3</sup> The Seven Countries study used 7-day weighed diet in 16 cohorts of subsets of men, and it only examined diet ecologically. Keys said “it would be prohibitively expensive and wholly unfeasible” to attain enough repeated measures to estimate true individual participant means of habitual diets;<sup>4</sup> individual practical Food Frequencies Questionnaires (FFQs) were not yet available 60 years ago, when the Seven Countries study was initiated. The PURE study<sup>1</sup> used FFQs in participants of both sexes and included cohorts within countries that were diverse enough to study socioeconomic status. Keys<sup>5</sup> selected countries to maximise a priori between-country differences in diet and coronary heart disease, whereas PURE<sup>1</sup> selected countries to maximise and represent countries with a broad range of gross national product per capita values. Both studies<sup>1,5</sup> had the practical restriction of requiring willing and competent investigators in the target countries.

In their Article in *The Lancet Global Health*, Annika Rosengren and colleagues<sup>6</sup> used PURE data to study individual educational attainment relative to incident cardiovascular disease and total death. The main finding was that low individual educational attainment predicted higher incidence of both outcomes, and disproportionately so in low-income countries. This finding was despite a relatively low prevalence of hypertension and diabetes in these people, but this result was also accompanied by high 28-day case fatality rates. Rosengren and colleagues<sup>6</sup> concluded that poor access to health care is the dominant factor in these associations. However, the most common

forms of cardiovascular disease are almost universally linked to risk factors. It is natural to wonder whether the cardiovascular disease in low-income countries among individuals with low education has atypical pathology. The disease incidence ascertainment method, relying heavily on participant and family reporting, would admit such a difference in pathology. However, the low education group in low-income countries is not actually free of cardiovascular disease risk factors: the diet score is directly correlated with individual educational attainment, and there were enough people who smoked (28%), had hypertension (29%), or diabetes (8%) for typical cardiovascular disease to be concentrated in these people. The authors do not present risk factor or behaviour relationships with cardiovascular disease incidence, so it is hard to judge the extent to which cardiovascular disease in this subset of PURE might be atypical. Our suggested conclusion is tempered, emphasising poor health behaviours along with poor access to primary and secondary health care. This is an important distinction for policy making.

A note on the statistical methods in the PURE work is revealing. Rosengren and colleagues<sup>6</sup> estimated individual-level associations with reasonably aggregated ecological units, namely low-income, middle-income, and high-income countries, then looked for individual-by-ecological interactions. This approach is what unveiled differential education-by-cardiovascular-disease associations by ecological grouping. The Seven Countries study<sup>7</sup> used a similar strategy in assessing whether serum cholesterol had the same association with mortality from coronary heart disease across six country groupings. The figure in the Seven Countries study<sup>7</sup> shows the mortality rate associated with coronary heart disease on the y-axis, serum cholesterol concentration on the x-axis, and six separate line graphs plotted, each indicating individual participant-based rate of coronary heart disease for a quartile of serum cholesterol within a subset of the seven countries. The ecological associations (between countries) are then seen as vertical differences. These graphs, scanned horizontally, revealed an increase in mortality from coronary heart disease with serum cholesterol in all country groupings except for Japan (a result that was

Published Online  
April 23, 2019  
[http://dx.doi.org/10.1016/S2214-109X\(19\)30187-1](http://dx.doi.org/10.1016/S2214-109X(19)30187-1)  
See **Articles** page e748

ambiguous because almost no one in the Japanese cohort had high cholesterol) yet, looking vertically in the display between countries, there were differential mortality rates associated with coronary heart disease between each country-pair (eg, northern Europe and the USA had higher rates than did southern Europe within each group of cholesterol concentrations). This important nuance suggests that serum cholesterol is causally important, but other factors must also be operating, and the association was possibly very weak in Japan. Only the individual-by-ecological interaction would reveal this pattern of association.

It is very tempting and common in analysis of data gathered over ecological units to pool the data without reference to the ecological unit. Mahshid Dehghan and colleagues<sup>8</sup> used this approach in studying total fat and total carbohydrate intake in the PURE study. Among their conclusions were that “high carbohydrate intake was associated with higher risk of total mortality” and that “global dietary guidelines should be reconsidered in light of these findings”. These are inordinately strong statements. Additional care must be taken in several respects. First, ecological variables do not necessarily have the same meaning as the individual variables with the same name;<sup>9</sup> for an extreme example, in random population samples, female sex is present in 50% of people, but this finding reflects population structure, not related sex-specific biology such as sex hormones. Second, examination of total carbohydrate or total fat ignores the effects of individual fatty acids and carbohydrates and dietary quality in general. Application of the methods of Rosengren and colleagues<sup>6</sup> or Verschuren and colleagues,<sup>7</sup> and addition of a general diet pattern score (such as was used in the study by Rosengren

and colleagues), has the potential to provide a clearer answer, addressing total carbohydrate in the context of how the rest of the diet might differ among countries. This approach might merit consideration in actions that depend on the truth of underlying causal assumptions, such as in formulating global dietary guidelines.

*\*David R Jacobs Jr, Daan Kromhout*

Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, Minneapolis, MN 55454, USA (DRJr); Department of Epidemiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands (DK)

jacob004@umn.edu

DRJr reports personal fees from California Walnut Commission. DK declares no competing interests.

Copyright © 2019 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

- 1 Teo K, Chow CK, Vaz M, Rangarajan S, Yusuf S. The Prospective Urban Rural Epidemiology (PURE) study: examining the impact of societal influences on chronic noncommunicable diseases in low-, middle-, and high-income countries. *Am Heart J* 2009; **158**: 1–7.
- 2 Keys A. Seven Countries—a multivariate analysis of death and coronary heart disease. Cambridge, MA: Harvard University Press, 1980.
- 3 Keys A, Menotti A, Karvonen MJ, et al. The diet and 15-year death rate in the Seven Countries study. *Am J Epidemiol* 1986; **124**: 903–15.
- 4 Keys A. The authors reply. *Am J Epidemiol* 1988; **128**: 239–41.
- 5 Pett KD, Willett WC, Vartiainen E, Katz DL. The Seven Countries study. *Eur Heart J* 2017; **38**: 3119–21.
- 6 Rosengren A, Smyth A, Rangarajan S, et al. Socioeconomic status and risk of cardiovascular disease in 20 low-income, middle-income, and high-income countries: the Prospective Urban Rural Epidemiologic (PURE) study. *Lancet Glob Health* 2019; published online April 23. [http://dx.doi.org/10.1016/S2214-109X\(19\)30045-2](http://dx.doi.org/10.1016/S2214-109X(19)30045-2).
- 7 Verschuren WM, Jacobs DR, Bloemberg BP, et al. Serum total cholesterol and long-term coronary heart disease mortality in different cultures. Twenty-five-year follow-up of the Seven Countries study. *JAMA* 1995; **274**: 131–36.
- 8 Dehghan M, Mente A, Zhang X, et al. Associations of fats and carbohydrate intake with cardiovascular disease and mortality in 18 countries from five continents (PURE): a prospective cohort study. *Lancet* 2018; **390**: 2050–62.
- 9 Jacobs DR Jr, McGovern PG, Blackburn H. The US decline in stroke mortality: what does ecological analysis tell us? *Am J Public Health* 1992; **82**: 1596–99.